

Quantum Liquid Signatures in Dwarf Stars

Gregory Gabadadze and David Pirtskhalava

*Center for Cosmology and Particle Physics, Department of Physics,
New York University, New York, NY 10003, USA*

Abstract

We develop further the proposal of arXiv:0806.3692 that a new state of matter – charged condensate of spin-0 nuclei – may exist in helium-core dwarf stars. The charged condensate and its fluctuations are described by an effective field theory Lagrangian. The spectrum of bosonic fluctuations is gapped, while electrons, at temperatures of interest, give rise to gapless excitations near the Fermi surface. These properties determine the evolution of the dwarfs with condensed cores. In particular, we show that such dwarf stars would cool significantly faster than their crystallized counterparts. As a result, the luminosity function for the helium-core dwarfs will have a sharp drop-off after the condensation. It is tempting to interpret the recently discovered abrupt termination of a sequence of 24 helium-core dwarf candidates in NGC 6397 as a signature of the charged condensation.

1. Introduction and Summary

There is roughly five orders of magnitude between what may be called the atomic scale, $Angström \sim 10^{-8} \text{ cm}$, and the nuclear scale, $Fermi \sim 10^{-13} \text{ cm}$. If in a neutral system of a large number of electrons and nuclei average inter-particle separations are between these two scales, then neither atomic nor nuclear effects will play any significant role. Moreover, the nuclei can be treated as point-like particles.

Such a system of electrons and nuclei constitutes cores of white dwarf stars. Up to a factor of a few, these are roughly Earth's size solar-mass objects; their central mass density may range over $\sim (10^6 - 10^{10}) \text{ g/cm}^3$, most of them being near the lower edge of this interval. Since the dwarf stars exhausted thermonuclear fuel in their cores already, they evolve by cooling [1]; the ones that we consider in this work cool from $\sim 10^8 \text{ K}$ down to lower temperatures.

At temperatures of interest the electron de Broglie wavelengths overlap with each other, and they obey the laws of quantum statistics. Hence, the electron properties are universal throughout all the dwarf stars; their number density may range in the interval $\sim (0.1 - 5 \text{ MeV})^3$, and at temperatures of interest they form a degenerate quantum gas (their Fermi energy is greater than their interparticle Coulomb interaction energy).

As to the nuclei – since they're heavier and their de Broglie wavelengths are shorter – they may exhibit different properties depending on the values of density and temperature at hand: they may crystallize [2] as it should be the case for a majority of dwarf stars, or alternatively, may form a quantum liquid [3], when densities are high enough. Identification and qualitative description of quantum liquid signatures in the cores of high-density cool white dwarf stars is a main subject of the present work.

As a typical dwarf star cools down, the Coulomb interaction energy in a classical plasma of charged nuclei will significantly exceed their classical thermal energy, and the nuclei, in order to minimize energy, would organize themselves into a crystal lattice [2]. In most of these cases quantum effects of the nuclei should be negligible; for instance, the Debye temperature should be less than the temperature at which crystallization takes place, and the de Broglie wavelengths of the nuclei should be much smaller than the average internuclear separations. This indeed is the case in majority of white dwarf stars, the cores of which are composed of carbon and/or oxygen nuclei and span the interval of mass densities around $\sim (10^6 - 10^8) \text{ g/cm}^3$.

However, there exists a class of dwarf stars in which the nuclei enter the quantum regime before the above-described classical crystallization process sets in [4, 5]. Among these, furthermore, there is a relatively small subclass of the dwarf stars for which the temperature T_c , at which the de Broglie wavelengths of the nuclei begin to overlap, is higher than the would-be crystallization temperature T_{cryst} (see [3] and discussions in Section 2 below). Then, right below T_c , the quantum-mechanical uncertainty in the position of the charged nuclei is greater than the average internuclear separation. This is diametrically opposite to the crystallized state where the

nuclei would have well-localized positions with slight quantum-mechanical fuzziness due to their zero-point oscillations.

What is then an adequate description of such a state? It was argued in [3, 6] that such a system, instead of forming a crystalline lattice at T_{cryst} , could condense at $T_c > T_{cryst}$, owing to the quantum-mechanical probabilistic “attraction” of Bose particles to occupy one and the same zero-momentum state, and leading to a quantum liquid in which the charged spin-0 nuclei would form a macroscopic quantum state with a large occupation number – the charged condensate.

It was proposed in [3] that the conditions in certain high-density helium-core white dwarfs (He WDs) are appropriate to form such charged condensate. Here we will argue that similar effects could take place in lower-density He WDs, which are more relevant for observations [7, 8, 9], as well as in superdense ($\sim 10^{10} \text{ g/cm}^3$) carbon-core white dwarfs (C WDs).

The observational consequences of the charged condensate in astrophysical objects can be significant. The bosonic part of the liquid is superconducting as its spectrum of long-wavelength fluctuations exhibits a mass gap; hence these quasi-particles contribute to the specific heat of the substance only in an exponentially suppressed way [3]. This affects the cooling rate for dwarf stars. In particular, we will show that the He WDs with the condensed cores would cool much faster than the ones in which condensation does not take place.

We should emphasize though that a careful consideration of white dwarf cooling rates requires detailed account of composition and dynamics of their envelopes, which is a complex and less certainly known subject (for detailed discussions see, e.g., [10, 11]). We will not attempt here to enter these studies. Instead, we consider an over-simplified model just to emphasize our main point that if the charged condensation takes place in the cores of white dwarfs, they’ll necessarily cool faster, and there will be a drop-off in their luminosity function.

One may wonder whether the effects that we’re discussing can be entirely obfuscated by the uncertainties in the envelope composition. Although this may well be the case for cooling times, there is nevertheless a model independent prediction of the charged condensation: the luminosity function will have a sharp drop-off after the condensation, with subsequent growth governed by a shallower slope.

The recently discovered and studied in Ref. [9] 24 helium-core dwarf candidates in NGC 6397 exhibit a termination of the sequence at low luminosities. It is tempting to attribute this to the drop-off due to the charged condensation. Whether this proposal can withstand more detailed scrutiny remains to be seen.

The organization of the paper is as follows: In Section 2 we give qualitative arguments why condensation may be preferred over crystallization in certain WDs. In Section 3 we summarize and further develop the effective field theory approach of [3, 12] to charged condensation, and describe the spectrum that is relevant for cooling of WDs. In Section 4 we calculate cooling times for He WDs and compare them with those of core-crystallized WDs. Last but not least, we discuss the drop-off of the luminosity function after charged condensation.

2. Condensation versus Crystallization

We concentrate on WDs that below certain temperature cool by releasing stored in them heat from their surface. At the beginning of that stage the ions (nuclei) form a classical Bose gas that cools with constant specific heat (the so-called Mestel cooling [1]) down to some temperature T_{cryst} , at which Coulomb repulsion starts to dominate by about two orders of magnitude over the average thermal energy, and the core undergoes crystallization transition [2]. The crystallization temperature in the classical regime is quantified by the ratio

$$\Gamma \equiv \frac{E_{\text{Coulomb}}}{2E_{\text{Thermal}}/3} = \frac{(Ze)^2/4\pi d}{k_B T}, \quad (1)$$

where e is the electric charge, Ze is the charge of a nucleus, k_B denotes the Boltzmann constant, J_0 is the electron number-density, $d \equiv (3Z/4\pi J_0)^{1/3}$ is the average inter-ion separation, while T denotes the core temperature of the star. The ion plasma is expected to undergo crystallization once the temperature drops to a value for which $\Gamma \simeq 180$ [13, 14, 15].

The above arguments are entirely *classical*. The temperature scale that determines the classical versus quantum nature of the crystallization transition is the Debye temperature

$$\theta_D \equiv \frac{\hbar \Omega_p}{k_B}, \quad \Omega_p = \left(\frac{J_H (Ze)^2}{m_H} \right)^{\frac{1}{2}}, \quad (2)$$

where Ω_p is the plasma frequency of the ion gas, $J_H = J_0/Z$ denotes the ion number density and m_H denotes the mass of a single ion (the subscript H stands for “heavy”). Up to a factor of $\sqrt{3}$, Ω_p coincides with the frequency of zero-point oscillations of the ions in crystal sites, $w_0 = \Omega_p/\sqrt{3}$.

The white dwarfs with $T_{\text{cryst}} > \theta_D$ cool according to the above-described classical scenario. Often however, θ_D may significantly exceed T_{cryst} [4]. In such a case, quantum zero-point oscillations should be taken into account in order to derive the crystallization temperature. This seems to delay the formation of quantum crystal, lowering T_{cryst} from its classical value at most by about $\sim 10\%$ [5]. Since this is a small change for the estimates that we’re after here, we will consider the classical value of T_{cryst} to be a good approximation, even in the quantum case, keeping in mind that T_{cryst} may overestimate somewhat the crystallization temperature of the substance.

However, there exists a third and very important temperature scale, relevant for studying the cooling of white dwarf interiors. It is the “critical” temperature T_c , at which the de Broglie wavelengths of the ions start to overlap

$$T_c \simeq \frac{4\pi^2}{3m_H d^2}. \quad (3)$$

Below T_c quantum-mechanical uncertainties in the ion positions become greater than an average inter-ion separation. Hence the latter concept loses its meaning as a microscopic characteristic of the system, and the ions enter a quantum-mechanical regime of indistinguishability¹. Below T_c the wavefunction of the many-body system of spin-0 ions should be symmetrized, and this would unavoidably lead to probabilistic “attraction” of the bosons to occupy the same quantum state.

Therefore, when crystallization temperature T_{cryst} is lower than T_c , the system may instead undergo condensation into a macroscopic zero-momentum quantum state with large occupation number – the charged condensate.

Once in the condensate, the boson positions are entirely uncertain while their momenta equal to zero. In order for such a system to crystallize later on, each of the bosons should acquire the momentum determined by the zero-point energy of the crystal ions, $\mathbf{k}_0^2 = 2m_H\Omega_p/\sqrt{3}$. The latter is greater than the Fermi momentum, as well as typical momenta of fermionic quasiparticles. Hence, the fermions will not be able to transmit to the condensed bosons momenta comparable with $|\mathbf{k}_0|$. Therefore, the transition to the crystallized state can only happen spontaneously. Such a transition could take place as it would lower the energy of the entire system due to the favorable electrostatic screening. However, given that the spectrum of bosonic quasiparticles is gapped, the process will be exponentially suppressed at temperatures below the gap scale. Hence, the condensate should not be expected to undergo subsequent crystallization, at least for a long period of time.

Another crucial question is whether the expression for the critical temperature (3) gives an accurate estimate for the actual condensation temperature T_{cond} at which the phase transition would take place. If we were to deal with a *non-interacting* system of Bosons then the known BE condensation temperature, $T_{\text{cond}}^{BE} \simeq (1.27/m_H d^2)$, would have been an order of magnitude smaller than what the estimate (3) suggests. However, it has been known that already *weak repulsive* interactions between bosons increase the condensation temperature; this is consistent with ones expectation that the repulsion makes easier for the condensation in the momentum space to take place (indeed, the BE condensation is a condensation in the momentum space, while the coordinate space wave-functions are entirely delocalized) see, e.g., [16] and references therein.

In the case of weakly interacting bosons the increase of the condensation temperature is small since the interactions are weak. In our case, however, interactions between spin-0 nuclei are strong, in a sense of a many-body system, as we are about to argue below. In this case we would expect $T_{\text{cond}} \gg T_{\text{cond}}^{BE}$. Since we have no means to evaluate T_{cond} accurately, we use the expression (3) as a reasonable and physically motivated approximation for the interacting system $T_c \sim T_{\text{cond}}$. To this end, the

¹The de Broglie wavelength above is defined as $\lambda_{dB} = 2\pi/|\mathbf{k}|$, where $\mathbf{k}^2/2m_H = 3k_B T/2$. We define T_c as the temperature at which $\lambda_{dB} \simeq d$. Note that this differs by a numerical factor from the standard definition of the *thermal* de Broglie wavelength, $\Lambda \equiv \sqrt{2\pi/mk_B T}$, that appears as a natural scale in the partition function. See comments on the rationale for our choice of T_c below.

criterion that we adopt for the condensation to take place is

$$T_c \gtrsim (a \text{ few}) T_{cryst}. \quad (4)$$

The charged condensate is somewhat similar, but also differs by its strong coupling, from the Bose-Einstein condensate of charged spin-0 particles. Analytic studies of a BE condensate of charged scalars in the past (see, e.g., Ref. [17] and references citing it) relied on a small departure from the condensation of free particles, and used an expansion in a parameter r_s that is defined as the ratio of the average interparticle separation to the would-be Bohr radius for the boson. For a weakly non-ideal system of bosons the r_s -ratio is small $r_s \ll 1$, and the ground state is the weakly-non-ideal BE condensate. Furthermore, as zero-temperature numerical simulations show (see, e.g., [18, 19], and references therein), for $1 \lesssim r_s \lesssim 160$ the condensate and crystal state begin to coexist. Furthermore, for $r_s \gtrsim 160$ the entirely crystallized state is a ground state. Presumably, the finite temperature effects would increase the value of r_s at which the crystallization takes place only by a factor of a few, but not by an order of magnitude (see, e.g., [20]).

In the system of nuclei and electrons the r_s -ratio cannot be made small without entering the regime where nuclear interactions become dominant. Instead, in the case considered in [3], $r_s \sim (10 - 100)$. In the present work we will consider even larger values, $r_s \sim 10^3$, which are relevant for observed He WDs. Yet we argue that because the temperature T_c at which the de Broglie wavelengths of bosons start to overlap is greater than the crystallization temperature, the system, upon gradual cooling, should settle in the charged condensate state due to the quantum-mechanical probabilistic properties of the indistinguishable bosons. In fact, our criterion $T_c > T_{cryst}$ implies that the charged condensation could take place at fine temperature T_c for $r_s < 2400$. It is also worth pointing out in this regard that the expansion parameter in our case ends up being $1/r_s$, as long as the system is described in an effective Lagrangian approach.

The seeming contradiction with the numerical results may be reconciled by the fact that the charged condensate can only be a metastable state [21], while the crystal should arguably be the lowest energy state for $r_s \gg 160$. Most of the numerical simulations use the test wavefunction approach that minimizes the energy; finding a long-lived state which represents only a local minimum may not be easy in this approach².

One check of this proposal is that the small fluctuations of the charged condensate have no unstable modes, suggesting that it represents at least a local minimum. Also, these fluctuations are rather different from those of a would-be crystal, ordinary clod plasma, or weakly-coupled BE condensate. For discussions of non-perturbative (meta)stability of the charged condensate in a different context, see [21].

²It is interesting to note that in Ref. [18] a metastable quantum liquid branch was observed for $r_s \gtrsim 160$. At this stage it is hard to speculate whether this is the branch that we're discussing here.

3. Charged Condensate and its Fluctuations

We use an effective Lagrangian description to study the charged condensate³. We focus on the zero-temperature limit, even though realistic temperatures in He WDs are well above zero (for calculations of the finite temperature effects, see [22]). The validity of the zero-temperature approximation is justified *a posteriori* and goes as follows: the spin-0 nuclei undergo the condensation to the zero-momentum state; while they do so they cannot excite their own phonons since the latter are gapped with the magnitude of the gap being greater than the condensation temperature. On the other hand, the condensing charged bosons can and will excite thermal fluctuations in the fermionic sector that is gap-less. Therefore, all the thermal fluctuations will end up being stored in the fermionic quasiparticles near the Fermi surface. For the later, however, the finite temperature effects aren't significant since their Fermi energy is so much higher, $T/J_0^{1/3} \ll 10^{-2}$.

After these comments we turn to the effective Lagrangian. By m_H we denote the mass of a (heavy) nucleus of charge Ze and atomic number A (helium-4, carbon, oxygen), by μ_f the electron chemical potential, and by m_e the electron mass. The following hierarchy of scales, $m_H \gg \max[\mu_f, m_e]$, is a starting point for the effective Lagrangian construction.

We begin at scales that are well below the heavy mass scale m_H , but somewhat above the scale set by $\max[\mu_f, m_e]$. Hence the electrons are described by their Dirac Lagrangian, while for the description of the nuclei we will use a charged scalar *order parameter* $\Phi(x)$. As it was shown in [12], in a non-relativistic approximation for the nuclei, an effective Lagrangian proposed by Greiter, Wilczek and Witten (GWW) [23] in a context of superconductivity, is also applicable for the description of the charged condensation, given that an appropriate reinterpretation of its variables and parameters is made.

The construction of the GWW effective Lagrangian is based on the following fundamental principles: it is consistent with the translational, rotational, Galilean and the global $U(1)$ symmetries, preserves the algebraic relation between the charged current density and momentum density, gives the Schrödinger equation for the order parameter in the lowest order, and is gauge invariant [23]. Combined with the electron dynamics the GWW effective Lagrangian reads (we omit for simplicity the standard Maxwell term):

$$\mathcal{L}_{eff} = \mathcal{P} \left(\frac{i}{2} (\Phi^* D_0 \Phi - (D_0 \Phi)^* \Phi) - \frac{|D_j \Phi|^2}{2m_H} \right) + \bar{\psi} (i\gamma^\mu D_\mu^f - m_f) \psi, \quad (5)$$

where we use the standard notations for covariant derivatives with the appropriate charge assignments: $D_0 \equiv (\partial_0 - iZeA_0)$, $D_j \equiv (\partial_j - iZeA_j)$, $D_\mu^f = \partial_\mu + ieA_\mu$, while

³An effective field theory that describes the charged condensate was discussed in Refs. [3, 6]. In this section we briefly describe and expand some of the results of these works.

$\mathcal{P}(x)$ stands for a general polynomial function of its argument ⁴.

The coefficients of this polynomial, $\mathcal{P}(x) = \sum_{n=0}^{\infty} c_n (x^n / \Lambda^{4n})$, are dimensionful numbers that are inversely proportional to powers of a short-distance cutoff of the effective field theory.

Once the basic Lagrangian is fixed, we introduce the electron chemical potential term $\mu_f \psi^+ \psi$, where $\mu_f = \epsilon_F = [(3\pi^2 J_0)^{2/3} + m_f^2]^{1/2}$. This gives a nonzero electron number density J_0 which is related to the Fermi momentum $k_F = (3\pi^2 J_0)^{1/3}$. This is also the only term that at the tree level sets a frame in which the electron total momentum is zero. The quantum loop corrections due to this term will generate additional Lorentz-violating terms in the bosonic sector of the theory [3, 22, 12].

There exists a solution of the equations of motion that follow from the effective Lagrangian (5). This solution takes the form [3]:

$$Z|\Phi|^2 = J_0, \quad A_\mu = 0, \quad \mathcal{P}'(0) = 1. \quad (6)$$

(We use the unitary gauge $\Phi = |\Phi|$). The condition $\mathcal{P}'(0) = 1$ is satisfied by any polynomial functions $\mathcal{P}(x)$ for which the first coefficient is normalized to unity

$$\mathcal{P}(x) = x + C_2 x^2 + \dots \quad (7)$$

The above solution describes a neutral system of negatively charged electrons of charge density $-eJ_0$, and positively charged scalar (helium-4 nuclei) condensate of charge density $Ze\Phi^+\Phi = eJ_0$. This describes the condensate and not a standard crystal in a long wavelength approximation where lattice inhomogeneities can be neglected, or ordinary cold plasma. This becomes more clear after one calculates the spectrum of small perturbations about the homogeneous solution and finds that it is rather different from the spectra of nearly free BE condensate, crystal lattice vibrations, or from plasma fluctuations.

Calculation of the spectrum of small perturbations is straightforward (here we follow conventions of [12]). There are two transverse polarizations of a massive photon which propagate with the conventional massive dispersion relation

$$\omega^2 = \mathbf{k}^2 + m_\gamma^2, \quad m_\gamma^2 \equiv \frac{Ze^2 J_0}{m_H}. \quad (8)$$

⁴In a more complete treatment one should also add to the Lagrangian terms $\mu_{NR} \Phi^* \Phi$, $\lambda(\Phi^* \Phi)^2 / m_H^2$, and the higher dimensional operators that are consistent with all the symmetries and conditions that lead to (5). Here μ_{NR} denotes a non-relativistic chemical potential for the scalars. These terms will not be important for the low-temperature spectrum of small perturbations we're interested in, as long as $\lambda \lesssim 1$ and $J_0 \ll m_H^3$. However, near the phase transition point it is the sign of μ_{NR} that would distinguish between the broken and symmetric phases, so these terms should be included for the discussion of the symmetry restoration. We also note that the scalar part of (5) is somewhat similar to the Ginzburg-Landau (GL) Lagrangian for superconductivity. However, there is a significant difference between the two. The coherence length in the GL theory is many orders of magnitude greater than the average interelectron separation, while in the present case, the “size of the scalar” Φ is smaller than the average interparticle distance.

Moreover, there is one longitudinal mode (phonon) with the following unconventional dispersion relation

$$\omega^2 \simeq m_\gamma^2 \left[1 + \frac{\mathbf{k}^4}{\mathbf{k}^2 m_0^2 + 4M^4} \right] \simeq m_\gamma^2 + \frac{\mathbf{k}^4}{4m_H^2}. \quad (9)$$

Here $M \equiv \sqrt{m_H m_\gamma}$, $m_0^2 \equiv m_\gamma^2 + \delta m^2$, where $\delta m^2 \propto e^2 J_0 / E_F$ is the Debye mass squared. As we see, $\omega^2(|\mathbf{k}| = 0) = m_\gamma^2$, and the bosonic collective excitations are gapped. Hence, the bosonic part of the whole system represents a superconducting component, while the fermions, at temperatures of interest, can be regarded as the “normal” (dissipative) component of the quantum liquid (see more below).

We point out that in the approximation used above there does not appear to be a $\mathcal{O}(\mathbf{k}^2)$ term on the r.h.s. of the dispersion relation (9). Such a term emerges (see [12]) in the $1/m_H^3$ order of the heavy mass expansion, and is proportional to $-m_\gamma^2 \mathbf{k}^2 / m_H^2$. This term is responsible for the “roton-like” behavior of small fluctuations discussed in [12]. The dispersion relation (9) is applicable for $k^2 \gtrsim m_\gamma^2$, which corresponds to distances shorter than $1/m_\gamma$ – the scale that encompasses a huge number of particles.

Note also that the phonon dispersion relation (9) is different from that of a conventional phonon in a crystal for which $\omega \propto |\mathbf{k}|$, near the origin. Hence, the substance that we’re discussing is different from a crystal, and it is also different from classical plasma. The following line of arguments also suggests that the charged condensate differs from the ordinary BE condensate of nearly-free bosons too: In the limit of switched off interactions, $e \rightarrow 0$, the dispersion relation (9) reduces to $\omega = \mathbf{k}^2 / 2m_H$. This is nothing but a dispersion relation for a lowest excitation in a BE condensate of *free* bosons of mass m_H . However, for realistic values of the parameters the second (momentum dependent) term in (9) is sub-dominant, suggesting that the charged condensate has a significant departure from the BE condensate of nearly-free bosons.

What is important for the present work is that the bosonic collective excitations give rise to exponentially suppressed contributions to the value of specific heat of this substance. Typical suppression scales as $\exp(-m_\gamma/T)$, and since for the He WDs $m_\gamma \simeq 3 \text{ KeV}$, at temperatures below 10^6 K , these contributions are proportional to $\sim \exp(-30)$ and can be neglected.

This is not the case in dwarf stars in which the nuclei formed a crystalline structure. There, the dominant contribution to the specific heat comes from a crystal phonon. The latter has a linear dispersion relation $\omega \propto |\mathbf{k}|$, and its contribution to the specific heat scales with temperature as T^3 .

Therefore, to a good approximation, there is no phonon contribution in the charged condensate case, while it is present when the cores crystallize.

As to the electrons, their behavior is similar in both crystal and condensate cases. At temperatures of interest they form a degenerate Fermi gas with gap-less excitations near the Fermi surface. Their contribution to the specific heat scales linearly with temperature. In the case of crystallized cores this is sub-dominant to the specific heat due to the crystal phonon. For the charged condensate, on the

other hand, the fermions are the dominant contributors to the specific heat. These properties make a difference in cooling of dwarf stars, as we'll discuss in detail in the next section⁵.

Finally, let us discuss briefly the question of impurity (hydrogen, helium-3, etc.) nuclei that may be present in the cores of white dwarfs. The static potential between two impurity nuclei of charge Q_1 and Q_2 consists of two parts [12]:

$$V_{stat} = \alpha_{em} Q_1 Q_2 \left(\frac{e^{-Mr}}{r} \cos(Mr) + \frac{4\alpha_{em}}{\pi} \frac{k_F^5 \sin(2k_F r)}{M^8 r^4} \right). \quad (10)$$

The first, exponentially suppressed term modulated by a periodic function, is due to cancellation between the screened Coulomb potential and that of a phonon. The second term, which exhibits a power-like behavior modulated by a periodic function, is due to the existence of gap-less excitations near the Fermi surface. This gives a generalization of the Friedel potential (see, e.g., [24]) to the case when in addition to the fermionic excitations the collective modes associated with the charged condensate are also taken into account⁶. As we see, the potential is not sign-definite. It can give rise to attraction between like charges; this attraction is due to collective excitations of both fermionic and bosonic degrees of freedom. This represents a generalization of the Kohn-Luttinger [26] effect to the case where on top of the fermionic excitations the collective modes of the charged condensate are also contributing (for the discussion of associated superconductivity, see footnote 5).

For the physical conditions present in dwarf stars, the second term in (10) is dominating. This term is strongly suppressed because of the phonon subtraction [12]. Thus, attraction between like charges due to this potential could give Cooper pairing between impurities (even if their concentration was significant enough) only at extremely small temperatures (see footnote 5).

⁵In the charged condensate Cooper pairs of electrons can also be formed, however, the corresponding transition temperature, and the magnitude of the gap, are suppressed by a factor $\exp(-1/e_{eff}^2)$, where e_{eff}^2 is proportional to the value of the inter-electron potential that contains both screened Coulomb and phonon exchange. The fact that this potential has attractive domain, but is very small, is suggested by the static potential found in [12] (see eq. (10) below); the latter is down by a power of a large scale M . In other words, the static potential and the zero-zero component of the propagator are both suppressed as $D_{00} \sim 1/r_s \sim (10^{-3} - 10^{-2})$, where r_s was discussed in Section 2. Moreover, taking into account the frequency dependence via the Eliashberg equation does not seem to change qualitatively the conclusion on a strong suppression of the pairing temperature.

Hence, even though the bosonic sector (condensed nuclei) is superconducting at reasonably high temperatures $\lesssim 10^6$ K, interactions with gap-less fermions could dissipate the superconducting currents. Only at extremely low temperatures, exponentially close to the absolute zero, the electrons could also form a gap leading to superconductivity of the whole system. In the present work we consider temperatures at which electrons are not condensed into Cooper pairs, and ignore the finite temperature effects.

⁶Note that for spin-dependent interactions the same effects of the charged condensate would give a generalization of the Ruderman-Kittel-Kasuya-Yosida (RKKY) potential [25].

3. White Dwarf Cooling

Most common dwarf stars have masses approximately equal to half of the mass of the Sun, central densities $\sim (10^6 - 10^8) \text{ g/cm}^3$ and are composed mainly of carbon and/or oxygen. Above the largest of the three temperature scales – T_{cryst} , θ_D and T_c – the cooling process is determined by thermodynamics of the classical Bose gas of the ions. For lower temperatures however, the state of the star is determined by the relative magnitude of these scales.

For present purposes, it is more convenient to rewrite the expressions for the temperature scales in terms of the *mass* density ρ measured in g/cm^3

$$T_{\text{cryst}} \simeq 10^3 \rho^{\frac{1}{3}} Z^{\frac{5}{3}} \text{ K}, \quad \theta_D \simeq 4 \cdot 10^3 \rho^{\frac{1}{2}} \text{ K}, \quad T_c = 3.5 \cdot 10^2 \rho^{\frac{2}{3}} / Z^{\frac{5}{3}} \text{ K}, \quad (11)$$

where the baryon number of an ion was assumed to equal twice the number of protons ($A = 2Z$) and $\Gamma \simeq 180$ was set.

Helium White Dwarfs

White dwarfs composed of helium constitute a smaller sub-class of dwarf stars (see, [9, 8] are references therein). They exhibit best conditions for the charged condensation. Most of helium dwarfs are believed to be formed in binary systems, where the removal of the envelope off the dwarf progenitor red giant by its binary companion happened before helium ignition, producing a remnant that evolves to a white dwarf with a helium core. Other astrophysical mechanisms for formation of isolated helium white dwarfs may also be possible [27]. In any event, helium dwarf masses range from $\sim 0.5 M_\odot$ down to as low as $(0.18 - 0.19) M_\odot$, while their envelopes are mainly composed of hydrogen. In this work we'll only consider He WDs whose hydrogen envelopes are thin, where no thermonuclear reactions are taking place. Such WDs cool by radiating off the stored in them heat.

Following [7], we will consider an over-simplified model of white dwarf cooling. Our treatment of what actually is an involved process, with significant uncertainties due to the envelope composition and opacity, should not be expected to give quantitatively precise predictions. Nevertheless, our approach is good enough, as we'll see, to capture the main difference that arises in cooling of dwarfs with the condensed cores.

For definiteness, we consider cooling of a reference helium star of mass $M = 0.5 M_\odot$ with the atmospheric mass fractions of the hydrogen, and heavy elements (metallicity) respectively equal to

$$X \simeq 0.99, \quad Z_m \simeq (0.0002 - 0.002). \quad (12)$$

The lower value of the metallicity $Z_m \simeq 0.0002$ is appropriate for the recently discovered 24 He WDs in NGC 6397 [9], but for completeness, we consider a wider

range for this parameter. Table 1 illustrates different physical characteristics of such a star with the mass density $\sim 10^6 \text{ g/cm}^3$.

The equation of state of a white dwarf interior is well modeled by the polytropes [7]. The average density of a non-relativistically degenerate star – which the helium dwarfs are an example of – may be about 5 times less than the central density. For the sake of simplicity, we will neglect the nontrivial density profile when dealing with thermodynamics of the dwarf stars and consider their average and central densities to be roughly equal. This won't lead to a significant error in the cooling analysis due to the fact that the variation of the mass density is not dramatic in the part of a star which encompasses most of its mass.

As seen from Table 1, critical temperature for the reference star significantly exceeds the crystallization temperature, while Debye temperature exceeds both. This star provides appropriate conditions for the formation of the charged condensate in its core.

Table 1: Values of physical quantities for a reference helium white dwarf

Physical quantity	Numerical value
Electron number density	$(0.13 \text{ MeV})^3$
Mass density	10^6 g/cm^3
Separation between atomic nuclei	10^3 fm
Crystallization temperature	$3 \cdot 10^5 \text{ K}$
Debye temperature	$4 \cdot 10^6 \text{ K}$
Critical temperature	10^6 K

In the Appendix we give a brief summary of the derivation of the luminosity-core temperature relation for white dwarfs (for details see, e.g., [7]). The definition of luminosity of a cooling star, combined with equation (29) of the Appendix, gives the expression for the cooling rate:

$$-c_v \frac{dT}{dt} = L = C A m_u T^{3.5}. \quad (13)$$

Here c_v denotes the specific heat per ion, m_u is the atomic mass unit and the constant C is inversely proportional to the atmospheric (envelope) metallicity of a star. In general, variations in atmospheric composition could change the cooling age significantly. Although the subject of atmospheric opacities is involved, the Kramer's approximation endowed with the atmospheric composition given in (12) is good enough for our purposes of comparing cooling rates of crystallized and condensed dwarfs.

The expressions for specific heats of different components of dwarf cores in corresponding regimes are given in Table 2.

To quantify the effects of the charged condensation on the cooling rate, we consider the ratio of cooling times for two identical helium dwarfs with and without the charged condensate in their interiors.

Table 2: Specific heats of different components of a white dwarf core

State	Specific heat
Classical Bose gas	$\frac{3}{2}k_B$
Quantum crystal	$\frac{16\pi^4}{5}\left(\frac{T}{\theta_D}\right)^3k_B$
Nonrelativistic Fermi Gas	$\frac{\pi^2}{2}\frac{k_B T}{E_F}k_B$
Relativistic Fermi Gas	$\frac{(3\pi^2)^{\frac{2}{3}}}{3}\frac{k_B T}{J_0^{1/3}}k_B$

Integrating (13), we find the following expression for the cooling time of a star in the Mestel regime

$$t_{He} = \frac{k_B}{CAm_u} \left[\frac{3}{5}(T_f^{-\frac{5}{2}} - T_0^{-\frac{5}{2}}) + Z \frac{\pi^2}{3} \frac{k_B}{E_F} (T_f^{-\frac{3}{2}} - T_0^{-\frac{3}{2}}) \right], \quad (14)$$

where T_f and T_0 denote the final and initial core temperatures. The first term in the bracket on the right hand side corresponds to cooling due to classical gas of the ions and the second term corresponds to the contribution coming from the nonrelativistic Fermi sea. The latter is sub-dominant in the range of final temperatures we are interested in (the factor Z in front of this term is due to Z electrons per ion). Since $T_f \ll T_0$, the age of a dwarf star typically doesn't depend on the initial temperature. Neglecting the fermion contribution, we find time that is needed to cool down to critical temperature $T_f = T_c$

$$t_{He} = \frac{3}{5} \frac{k_B T_c M}{Am_u L(T_c)} \simeq (0.76 - 7.6) \text{ Gyr}. \quad (15)$$

Where an order of magnitude interval in (15) is due to the interval in the envelope metallicity composition given in (12). We also find the corresponding luminosities

$$L(T_c) \simeq (10^8 \text{ erg/s}) \frac{M}{M_\odot} \left(\frac{T_c}{\text{K}} \right)^{7/2} \simeq 1.5 \cdot (10^{-4} - 10^{-5}) L_\odot, \quad (16)$$

which are in the range of observable luminosities ($L_\odot \simeq 3.84 \cdot 10^{33} \text{ erg/s}$).

After the condensation, specific heat of the system dramatically drops as the collective excitations of the condensed nuclei become massive and “get extinct”. A contribution from the Fermi sea, which is strongly suppressed by the value of Fermi energy, becomes the dominant one. The phase transition itself would take some time to complete, and the drop-off in specific heat will not be instantaneous. During that time a heat-transfer from the bosonic sector to the fermionic one will take place, but this will only change temperature of the fermions by a factor of $(1 + 1/Z)$, which, given our approximations, can be ignored. Effects of finite duration of the phase transition on the shape of the luminosity function will be discussed in the next section. Here, for simplicity we approximate the transition to be instantaneous, and

retain only the fermion contribution to specific heat below T_c . Then, the expression for the age of the star for $T_f < T_c$, reads as follows

$$t'_{He} = \frac{k_B}{CAm_u} \left[\frac{3}{5}(T_c^{-\frac{5}{2}} - T_0^{-\frac{5}{2}}) + Z \frac{\pi^2}{3} \frac{k_B}{E_F}(T_f^{-\frac{3}{2}} - T_0^{-\frac{3}{2}}) \right]. \quad (17)$$

Notice the difference of (17) from (14) – in the former $T_f < T_c$ and it is T_f that enters as final temperature in the fermionic part, while T_c should be taken as the final temperature in the bosonic part.




Figure 1: The ratio of the ages, as a function of final temperature, for two identical helium dwarf stars, with and without the interior condensation.

Figure 1 gives the ratio of ages, $\eta = t_{He}/t'_{He}$, for two identical helium dwarf stars, with and without the interior condensation. The considered range of final temperature is $(3 \cdot 10^5 - 10^6) K$. At the upper edge of this interval, the luminosity of the star, depending on the metallicity interval given in (12), is $(10^{-4} - 10^{-5})L_{\odot}$, while for the lower edge of the temperature interval we get $2 \cdot (10^{-6} - 10^{-7})L_{\odot}$. As is seen from this figure, the charged condensation substantially increases the rate of cooling. At the lower end of the considered temperature interval the age is about twenty times less than it would have been without the condensation phase.

With the decrease of density, the critical and Debye temperatures drop as $T_c \propto \rho^{\frac{2}{3}}$ and $\theta_D \propto \rho^{\frac{1}{2}}$ respectively, whereas the crystallization temperature drops only as $T_{cryst} \propto \rho^{\frac{1}{3}}$. Hence, the critical and crystallization temperatures would become equal for the dwarfs of low density. This would happen for densities $\sim 2 \cdot 10^4 g/cm^3$, which are very low and unlikely to be present in white dwarfs cores. Hence, we'd expect that all He WDs should have a large fractions of their cores in the condensed state.

To summarize, the charged condensation of white dwarf interiors would have a significant impact on cooling curves of helium dwarf stars at the faint end of the luminosity spectrum. Starting from a certain temperature scale, $\sim T_c$, the ages of the stars become effectively "frozen", whereas the temperature and luminosity decrease rather fast. Such a change in the cooling regime would leave its signatures on the luminosity function of helium white dwarfs, which we'll discuss below.

Luminosity Function

The luminosity function (LF) $\phi(\frac{L}{L_\odot})$ can be used to test the white dwarf evolution models against the observational data; it is defined as follows:

$$\phi(\frac{L}{L_\odot})d(\log(\frac{L}{L_\odot})) \equiv \text{number-density of WD's per unit interval of } (\log(\frac{L}{L_\odot})). \quad (18)$$

Under the assumption of the standard star formation rate, uniform both in space and time, the luminosity function takes a simple power-law form

$$\phi \propto \left[\frac{d \log(\frac{L}{L_\odot})}{dt} \right]^{-1} \propto L^{\frac{n(k+1)}{4}-1}, \quad (19)$$

where the constants k and n are the exponents defining temperature dependence of specific heat and luminosity:

$$c_v \propto T^k, \quad L \propto T^{\frac{4}{n}}. \quad (20)$$

For instance, n equals to $8/7$, once the Kramer's opacity is adopted for the description of the dwarf atmosphere. The log of the LF corresponding to cooling in the Mestel regime ($c_v = 3k_B/2$), is just a line with a slope equal to $5/7$.

The condensation of the core would induce significant deviations from the Mestel curve for helium white dwarfs. The exponential suppression of specific heat of the ions in the condensate doesn't occur instantaneously. Typically, for a quantum liquid, there is a transition region of width $\sim \Delta T_c$, in which the specific heat changes in a certain way. Therefore, starting from some temperature $T_c - \Delta T_c$, at which the specific heat of the ions can be neglected and a stage of rapid cooling begins (the age of the star "freezes"), the luminosity function starts to drop dramatically (Fig. 2). The stars, previously belonging to the same logarithmic luminosity bin, will spread among a few bins as a result of suddenly accelerated cooling.

Not much can be said with certainty of ΔT_c and the temperature dependence of specific heat near the phase transition point. In a well-known case of quantum liquid of neutral helium atoms for instance, the specific heat exhibits the so-called "Lambda" behavior. In analogy with the latter, one may expect the specific heat to increase near the phase transition point before dropping dramatically below T_c . If so, the parameter ΔT_c would give the width of the transitional region. Depending on the details of the function $c_v(T)$ near the phase transition point, the luminosity function




Figure 2: A schematic sketch of the luminosity function for helium-core white dwarfs. The absolute normalization of the function is set by the constant b which is determined by their formation rate. The blue line represents the Mestel regime. The shape of the luminosity function near the condensation point depends on the details of the corresponding phase transition.

of He WDs could take different shapes near its maximum, as it is schematically illustrated in Fig. 2.

What is independent of the above uncertainties, however, is the fact that the LF will experience a significant drop-off after the charged condensation phase transition is complete. This is due to the “extinction” of the bosonic quasiparticles below the phase transition point. In fact, the LF function will drop by a factor of ~ 200 , which would be reflected as a drop-off by about two units of the vertical axis from its maximum shown on Fig 2. After the drop-off the log of the LF function starts growing again but now with the slope equal to $3/7$ – due to the specific heat of the electrons. The latter part of the LF is shown only for one of the curves on Fig 2. Whether this drop-off can explain the termination of the 24 He WD sequence found in [9], remains to be seen in more detailed studies.

Superdense Carbon White Dwarfs

Finally, we briefly mention another possible subclass of white dwarf stars, that could undergo the core condensation. These are superdense dwarfs composed mainly of carbon, with masses close to the Chandrasekhar limit $\sim 1.4 M_\odot$, and central densities $\sim 10^{10} g/cm^3$. Such a star would be very close to the neutronization threshold ($5 \cdot 10^{10} g/cm^3$ for carbon nuclear matter), and its critical temperature $T_c \sim 8 \cdot 10^7 K$ would be greater than the crystallization temperature $T_{\text{cryst}} \sim 4 \cdot 10^7 K$, making condensation a possibility⁷.

Assuming the helium dominated envelope (Y denotes the helium fraction) with a 10% metallicity

$$X \simeq 0, \quad Y \simeq 0.9, \quad Z_m \simeq 0.1, \quad (21)$$

we find the cooling time needed to reach down to temperature $T_c \sim 8 \cdot 10^7 K$

$$t_C = \frac{3 k_B T_c M}{5 A m_u L} \simeq 3 \cdot 10^{-3} \text{ Gyr}. \quad (22)$$

This temperature corresponds to luminosity $L \simeq 3.4 L_\odot$. Once the star enters the condensation regime, it cools faster. For instance, it would reach the luminosity $L \simeq 10^{-6} L_\odot$ in $t = 0.7$ Gyr after the condensation. On the other hand, if one assumes that this star crystallizes instead of condensing its core, then it would reach the crystallization point in 10^{-2} Gyr, while it would take only ~ 0.1 Gyr longer for this star to cool down to $L \simeq 10^{-6} L_\odot$, as compared with the condensation case. In the crystallized phase for this star the cooling due to the fermion contributions is significant. Therefore, the log of the LF for such a star would not have a sharp

⁷At these high temperatures the WD cooling rate is significantly affected by the neutrino emission. Our goal, however, is to identify qualitative differences of cooling in the condensate and crystalline phases. For this purpose, and for simplicity, we ignore the neutrino contributions, which should certainly be taken into account for these WDs in more precise studies.

drop-off; instead, after the phase transition the LF would just change its slope to $3/7$ in the low luminosity region.

We also note that for dwarf stars with relativistically degenerate cores, the polytrope models yield more dramatic density profiles – density in the center exceeds the average density by a factor of ~ 50 . Since the crystallization/critical temperature equality is achieved at densities $\sim 10^9 \text{ g/cm}^3$ for carbon interiors, we should not expect the entire core to condense. In realistic calculations of cooling rates of the superdense carbon WDs this should also be taken into account.

Magnetic properties of condensed WDs, and their similarity to type II superconductors, will be discussed in [28].

Finally, we mention that charged condensation may also take place in some other astrophysical objects where the densities and temperatures are appropriate. Crusts of neutron stars may be a place to look at.

Acknowledgments

We'd like to thank Paul Chaikin, Daniel Eisenstein, Leonid Glazman, Andrei Gruzinov, Andrew MacFadyen, Aditi Mitra, Slava Mukhanov, Rachel Rosen and Malvin Ruderman for useful discussions and correspondence. GG was supported by the NSF and NASA grants PHY-0758032, NNG05GH34G. DP acknowledges the NYU James Arthur graduate fellowship support.

Appendix

For purposes of studying their cooling, white dwarfs are well described by a simple 2-component model. The core consists of the ion gas and degenerate electrons with large mean free path and heat conductivity, making it almost isothermal. The cooling takes place through the non-degenerate surface layer, the envelope, that surrounds the core. The photon diffusion equation, describing the energy flow from the core to the outer layers, has the following form

$$L = -4\pi r^2 \frac{c}{3\kappa\rho} \frac{d}{dr}(aT^4), \quad (23)$$

where L is luminosity, aT^4 - energy density of a blackbody, κ is the opacity of the stellar matter and ρ -mass density. We can rewrite the last equation in the following form

$$\frac{dT}{dr} = -\frac{3}{4ac} \frac{\kappa\rho}{T^3} \frac{L}{4\pi r^2}. \quad (24)$$

For the opacity, we use Kramer's approximation $\kappa = \kappa_0 \rho T^{-3.5}$.

The equation of the hydrostatic equilibrium (where P denotes the pressure and $m(r)$ - the mass inside a sphere of radius r)

$$\frac{dP}{dr} = -\frac{Gm(r)\rho}{r^2}, \quad (25)$$

combined with the equation of state for a classical ideal gas

$$P = \frac{\rho}{\mu m_u} k_B T, \quad (26)$$

(where μ and m_u are the mean molecular weight and atomic mass unit respectively), can be integrated with the boundary condition $P = 0$ at $T = 0$ to obtain

$$\rho = \left(\frac{2}{8.5} \frac{4ac}{3} \frac{4\pi GM}{\kappa_0 L} \frac{\mu m_u}{k_B} \right)^{\frac{1}{2}} T^{3.25}, \quad (27)$$

where $\kappa_0 = 4.34 \cdot 10^{24} Z_m(1 + X) \text{ cm}^2/g$ is the commonly used value for the opacity constant, X is the mass fraction of hydrogen, and Z_m - that of heavy elements (all elements except hydrogen and helium), while M is the mass of the star. Equation (27) works as long as we deal with nondegenerate matter. To estimate the limits of its applicability, we should equate the pressure of nondegenerate *electrons* (obtained from (26) by replacing μ by mean molecular weight per electron μ_e) to that of the nonrelativistic electron gas in the outer layers of the core

$$\frac{\rho_* k_B T_*}{\mu_e m_u} = \frac{3^{\frac{2}{3}} \pi^{\frac{4}{3}}}{5} \frac{\hbar^2 \rho_*^{5/3}}{m_e (m_u \mu_e)^{5/3}}, \quad (28)$$

where ρ_* and T_* are the density and temperature of the star at the core-surface boundary. Combining equations (28) and (27), we obtain a useful relation for the dependence of the luminosity on the temperature at the core-surface boundary (which, due to the isothermality of the core, can be used as a good characteristic of the core temperature)

$$L = \gamma \frac{\mu}{\mu_e^2} \frac{1}{Z_m(1 + X)} \frac{M}{M_\odot} (T_*)^{\frac{7}{2}} \equiv CM(T_*)^{\frac{7}{2}}, \quad (29)$$

where $\gamma \simeq 5.7 \cdot 10^5 \text{ erg/s}$ is a universal constant, and C is a constant that depends on the chemical composition of the envelope and varies from a star to star. A more general form of the last equation may be written as follows:

$$L \propto T_n^{\frac{4}{n}}, \quad (30)$$

where n is a constant, equal to $8/7$ in case of the envelope with Kramer's opacity.

References

- [1] L. Mestel, 1952, MNRAS, 112, 583.
- [2] L. Mestel and M.A. Ruderman, MNRAS, 136:27 (1967).
- [3] G. Gabadadze and R. A. Rosen, JCAP **0810**, 030 (2008) [arXiv:0806.3692 [astro-ph]].
- [4] G. Chabrier, N.W. Ashcroft, and H.E. DeWitt, Nature, 360,48 (1992)
- [5] G. Chabrier, Ap.J. 414:695 (1993).
- [6] G. Gabadadze and R. A. Rosen, Phys Lett. **B** 658 (2008), 266 [arXiv:0706.2304 [hep-th]].
- [7] S.L. Shapiro and S. A. Teukolsky, Black Holes, White Dwarfs, and Neutron Stars, John Wiley & Sons, (1983).
- [8] J. Liebert, P. Bergeron, D. Eisenstein, H.C. Harris, S.J. Kleinman, A. Nitta, J. Krzesinski, ApJ, **606**, L147, (2004), [astro-ph/040429].
- [9] R. R. Strickler, A.M. Cool, J. Anderson, H. N. Cohn, P. M. Lugger, A.M. Serenelli, arXiv:0904.3496, astro-phGA.
- [10] F. D’Antona, and I. Mazzitelli, Ann. Rev. A. A., **28**, 139 (1990)
- [11] B.M.S. Hansen, J. Liebert, Annu. Rev. Astron. Astrophys. **41**, 465. (2003)
- [12] G. Gabadadze and R. A. Rosen, JCAP **0902**, 016 (2009) [arXiv:0811.4423 [hep-th]].
- [13] D.Q. Lamb and H.M. Van Horn, Ap.J. **200**:306 (1975).
- [14] S. Ichimaru, Rev. Mod. Phys. **54**:1017 (1982).
- [15] G.S. Stringfellow, H.E. DeWitt, W.I. Slattery, Phys. Lett. **A** **41**, 1105 (1990).
- [16] K. Huang, Phys. Rev. Lett. **83**, 3770 (1999).
- [17] L.L. Foldy, Phys. Rev. **124**, 649 (1961).
- [18] D.M. Ceperlley, B.J. Alder, Phys. Rev. Lett. **45**, 566 (1980).
- [19] S. Moroni, S. Conti, M.P. Tosi, Phys. Rev. **B** **53**, 9688 (1996).
- [20] G. Chabrier, F. Douchin, A.Y. Potekhin, J. Phys.: Cond. Matter, **14** 9133 (2002).

- [21] G. Gabadadze and R. A. Rosen, Phys. Lett. B **666**, 277 (2008) [arXiv:0801.0228 [hep-th]].
- [22] A. D. Dolgov, A. Lepidi and G. Piccinelli, JCAP **0902**, 027 (2009) [arXiv:0811.4406 [hep-th]].
- [23] M. Greiter, F. Wilczek and E. Witten, Mod. Phys. Lett. B **3**, 903 (1989).
- [24] A.L. Fetter, J.D. Walecka, “*Quantum Theory of Many-Particle Systems*”, McGraw-Hill, 1971.
- [25] M.A. Ruderman and C. Kittel, Phys. Rev. **96**, 99 (1954); T. Kasuya, Prog. Theor. Phys. **16**, 45 (1956); K. Yosida, Phys. Rev. **106**, 893 (1957).
- [26] W. Kohn, J.H. Luttinger, Phys. Rev. Lett. **15**, 524 (1965).
- [27] M. Castellani, V. Castellani, ApJ., **407**, 649 (1993).
- [28] G. Gabadadze, R.A. Rosen, “Vortex Structure in Charged Condensate” NYU-TH-09/04/16; in preparation.